

# Independent components of the Riemann and Ricci tensors

Edoardo Milotti

October 20, 2022

The Riemann curvature tensor has 4 indexes, therefore it has 256 components. However not all the components are different, and here we count the actual number of different components.

Recall the identities

$$R_{\mu\alpha\beta\gamma} = -R_{\alpha\mu\beta\gamma} = -R_{\mu\alpha\gamma\beta} = R_{\beta\gamma\mu\alpha}$$

, (the Riemann tensor is antisymmetric with respect to the exchange of both the first pair of indexes and the second pair of indexes, while it does not change in a double exchange  $1 \leftrightarrow 3$ ,  $2 \leftrightarrow 4$ ) and

$$R_{\mu\alpha\beta\gamma} + R_{\mu\beta\gamma\alpha} + R_{\mu\gamma\alpha\beta} = 0$$

(the *cyclic identity*).

The first antisymmetry means that if  $\alpha = \mu$  then  $R_{\mu\alpha\beta\gamma} = 0$ . In order to obtain a nonzero value there are 4 choices for the first index and 3 for the second one. Since order does not matter, there are in all 6 different pairs. The same holds for the second pair of indexes. This means that there are potentially 36 nonzero independent components. However, the third equality  $R_{\mu\alpha\beta\gamma} = R_{\beta\gamma\mu\alpha}$  means that one can exchange the first pair with the second pair without altering the value of the coefficient (in practice, the matrix of values indexed by the  $\mu\alpha$  pair for the rows and the  $\beta\gamma$  pair for the columns is symmetrical), and therefore there are at most  $n(n+1)/2 = 6 \times 7/2 = 21$  independent values (recall that  $n = 6$  is the number of rows and columns of this matrix).

We still have to use the last identity

$$R_{\mu\alpha\beta\gamma} + R_{\mu\beta\gamma\alpha} + R_{\mu\gamma\alpha\beta} = 0,$$

where we can raise the first index to find

$$R_{\alpha\beta\gamma}^{\mu} + R_{\beta\gamma\alpha}^{\mu} + R_{\gamma\alpha\beta}^{\mu} = 0.$$

However, it is easy to see that whenever two indexes are equal, this is identically null simply because of the previous symmetry identities (Exercise: verify this statement). Therefore, the only additional constraint is

$$R_{0123} + R_{0231} + R_{0312} = 0$$

and this brings down to 20 the total number of independent components of the Riemann curvature tensor.

Now, recall that the Ricci tensor is defined by

$$R_{\mu\nu} = R_{\mu\nu\alpha}^{\alpha}$$

and note that by contracting the cyclic identity over  $\mu$  and  $\gamma$ , we find

$$R_{\alpha\beta\mu}^{\mu} + R_{\beta\mu\alpha}^{\mu} + R_{\mu\alpha\beta}^{\mu} = 0.$$

We know that all the components of the Riemann tensor with repeated indices vanish, therefore

$$R_{\alpha\beta\mu}^{\mu} + R_{\beta\mu\alpha}^{\mu} = 0,$$

and because of the antisymmetry of the last two indexes  $R_{\beta\mu\alpha}^{\mu} = -R_{\beta\alpha\mu}^{\mu}$ , and finally

$$R_{\alpha\beta\mu}^{\mu} = R_{\beta\alpha\mu}^{\mu},$$

which means

$$R_{\alpha\beta} = R_{\beta\alpha},$$

i.e., the Ricci tensor is symmetric and has only 10 independent components